

Brand Name: PITAS Cough Troche

RISK CATEGORIES

Second-class OTC drugs

CHARACTERISTICS

- This medicine contains oral disinfectant cetylpyridinium chloride hydrate to suppress sore throat and swelling caused by inflammation of the throat.
- The cough-repellent ingredient dextromethorphan phenolphthalinate acts in the cough center and suppresses cough.



ACTIVE INGREDIENTS (In 6 troches)

Dextromethorphan Phenolphthalinate 60 mg
Potassium Guaiacolsulfonate 140 mg Cetylpyridinium Chloride Hydrate 6 mg

[Inactive ingredients]: Hydroxypropylcellulose, Povidone, Pullulan, Macrogol, l-Menthol, Tannic Acid, D-Sorbitol, Sodium Saccharine, Sucrose Fatty Acid Ester, Sucralose, Sodium Copper Chlorophyllin, Flavor

For further information:

Manufactured by	TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form	Troche		
Packaging unit	12 troches		
Manufacturer's suggested retail price	JPY 886 (JPY 805 excluding tax)		
JAN Code	45198625		
Expiration date	3 years		

INDICATIONS

Cough
Phlegm
Throat pain, throat swelling, sore throat, throat discomfort, and hoarse voice due to throat inflammation

DOSAGE AND ADMINISTRATION

To include the following amounts in the mouth and use them by melting slowly first.

Age	Amount per time	Number of daily use	Service interval
Adults (15 years and older)	1 pc	6 times	2 hours or more
Under 15 years old	Do not use.		

<Notes Related to Dose and Administration>

- (1) Strictly observe the prescribed dose and administration.
- (2) Do not chew or stick in.

PRECAUTIONS FOR USE

The precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

1. The following persons should not take this medicine:

- Persons having a history of allergy to this product or any component of this product.

2. During use of this product, the following drugs may not be used.

Other antitussive expectorants, cold medicines, sedatives, oral-dose drugs containing antihistamine, and so on (oral-dose drugs for rhinitis, drugs for motion sickness, anti-allergy drugs, etc.)

3. Do not drive a car or operate machinery after taking this product.

(drowsiness may occur)

Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using this medicine.

- (1) Persons under treatment by a physician
- (2) Pregnant or possibly pregnant women
- (3) Persons having a history of allergy to drugs, etc.
- (4) Persons having any of the following symptoms: High fever.

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of this medicine immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected area	Skin	Gastrointestinal System	Neuropsychiatric System	Respiratory System
Symptoms	Eruption/redness, itching sensation	nausea/vomiting, anorexia	dizziness	dyspnea, shortness of breath

3. The following symptoms may be observed after taking this product. If these symptoms persist or worsen, discontinue the use of this product, and show this document to a doctor, pharmacist, or registered salesperson for a consultation

Drowsiness

4. If the symptoms do not improve after using the medicine for 5-6 times, discontinue the use of this medicine, and consult a physician, pharmacist or registered salesperson with this leaflet.

Precautions in storage and handling

- (1) Store in a cool place with little moisture, not exposed to direct sunlight.
- (2) Store in a place out of the reach of children.
- (3) Do not replace other containers (cause misuse or change quality).
- (4) Use immediately after opening an aluminum bag.
- (5) Do not use products that have passed their expiration date.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following: Distributor

Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27
Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66

Business hours: 9:00-17:00 (Excluding weekends and holidays) Website:

<https://www.taiho.co.jp/>