


Brand name: ZENOL XUM FX			
<b>RISK CATEGORIES</b> Second-class OTC drug			
<b>CHARACTERISTICS</b> <ul style="list-style-type: none"> <li>● Topical analgesic anti-inflammatory agent Felbinac (3%) helps relieve pains in shoulders, low back, etc.</li> </ul> As it is transparent and not sticky, it allows putting on clothes immediately after use. It is an easy-to-carry slim stick. It has a flavor of faint floral green.			
<b>ACTIVE INGREDIENTS</b> Per 100 g		For further information:	
		Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.
<b>ACTIVE INGREDIENTS</b>	Content	Manufactured and Distributed by:	Mikasa Seiyaku Co., Ltd.
Felbinac	3.0 g	Dosage form	Stick type agent
L-Menthol	2.7 g	Packaging unit	32 g x 1 box
Inactive ingredients: Macrogol, Crotamiton, Oleyl Alcohol, Stearic Acid, Isostearic Acid, Sodium Lactate, Flavor, Phenylethyl Alcohol, Squalane, pH Adjuster, Isopropanol, and other two components		Manufacturer's suggested retail price	JPY 1,455 (JPY 1,323 excluding tax)
		JAN Code	4987117372038
		Expiration	3 years
		<b>DOSAGE AND ADMINISTRATION</b> Apply an appropriate amount to the affected area twice to four times daily.  <b>&lt;Precautions&gt;</b> (1) Strictly follow the dosing instructions. (2) Be careful not to get the agent into the eyes. If the agent gets into the eyes, immediately wash with water or lukewarm water. If symptoms are severe, consult an ophthalmologist. (3) For external use only. Do not take the agent orally. (4) Do not cover the affected area with a poorly-ventilated material such as wrap film, etc. after applying the agent. (5) Wipe off sweat on the affected area before use. (6) Twist the container until 4 to 5 mm of agent appears before use, to prevent the container from directly touching the affected area. (7) Stop using the agent if the bottom of the container appears and becomes unstable. Do not forcibly use the agent remaining in the bottom of the container.	

<p><b>INDICATIONS</b> Shoulder pain due to stiff shoulder, low back pain, joint pain, muscle pain, bruise, sprain, tenosynovitis (pain and swelling in hands, wrists, and ankles), elbow pain (tennis elbow, etc.)</p>	
--	--

**PRECAUTIONS FOR USE**

☒ The following precaution should be observed.  
(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

1. The following persons should not use the agent.
  - (1) Persons with a history of allergic symptoms to the agent or any of the ingredients of the agent
  - (2) Persons with a history of asthma
  - (3) Pregnant women or possibly pregnant women
  - (4) Children under 15 years
2. Do not apply to the following areas.
  - (1) Around the eyes, mucous membranes, etc.
  - (2) Eczema, contact dermatitis, wounds
  - (3) Athlete's foot (tinea pedis), tinea cruris, etc. or purulent areas

■ Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using the agent.
  - (1) Persons under treatment by a physician
  - (2) Persons with a history of allergic symptoms to any drug
2. The following symptoms may be adverse reactions to the agent. If any of these symptoms occur, discontinue the use of the agent immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected body system	Symptoms
Skin	Rash, redness, itching, swelling, tingling sensation, contact dermatitis

The following serious symptom may rarely occur. If the symptom occurs, consult a physician immediately.

Symptom	Symptoms
Shock (anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, larynx itching, difficulty breathing, palpitation, consciousness clouding, etc. occur immediately after using the agent.

3. If symptoms do not improve after using the agent for about one week, discontinue the use of the agent, and consult a physician, pharmacist or registered salesperson with this leaflet.

**CONTACT**

For inquiries, please contact the shop where you purchased the product or the following:  
 Distributor Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.  
 1-27 Kandnishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan  
 Phone: 0120-4527-66  
 Business hours: 9:00-17:00 (Excluding weekends and holidays)  
 Website: <https://www.taiho.co.jp/>

Manufacturer and distributor : MIKASA SEIYAKU CO., LTD