


Brand name: Henseki																							
<p>RISK CATEGORIES Second-class OTC drug</p> <p>CHARACTERISTICS</p> <ul style="list-style-type: none"> This is a medicine to reduce adiposity It contains 11 types of crude drugs to reduce abdominal fat and excess fat in the body. Crude drugs contained in Henseki excrete waste products and water from stool and blood, and activate the metabolism. The medicine is suitable for persons with all constitutional types. Henseki is an originally formulated herbal medicine. As the medicine is suitable for persons with all constitutional types, it is recommended to reduce excess fat. No additives The medicine is manufactured only from crude drugs, and contains no additives. 																							
	<p>ACTIVE INGREDIENTS</p> <p>Three pouches (6 g) contain the following crude drugs.</p> <p>Powdered Alisma Tuber...0.86 g, Powdered Rhubarb...0.43 g Powdered Ginger...0.43 g, Powdered Glycyrrhiza...0.43 g Powdered Cinnamon Bark...0.43 g, Powdered Peony Root...0.43 g Powdered Moutan Bark...0.43 g, Powdered Polyporus Sclerotium...0.86 g Powdered Bupleurum Root...0.86 g, Powdered Pinelliae Tuber...0.43 g Powdered Cimicifuga Rhizome...0.43 g No additives</p> <p><Precautions> As this medicine is manufactured from crude drugs, slight difference in color and taste may occur between each product, but will not affect the quality.</p>	<p>For further information:</p> <table border="1"> <tr> <td>Distributed by:</td> <td colspan="2">TAIHO PHARMACEUTICAL CO., LTD.</td> </tr> <tr> <td>Manufactured and Distributed by:</td> <td colspan="2">Tatebayashi Shokakudo Co., Ltd.</td> </tr> <tr> <td>Dosage form</td> <td colspan="2">Powder</td> </tr> <tr> <td>Packaging unit</td> <td>2 g x 21 pouches</td> <td>2 g x 60 pouches</td> </tr> <tr> <td>Manufacturer's suggested retail price</td> <td>JPY 2,404 (JPY 2,185 excluding tax)</td> <td>JPY 6,072 (JPY 5,520 excluding tax)</td> </tr> <tr> <td>JAN Code</td> <td>4987117670448</td> <td>4987117670417</td> </tr> <tr> <td>Expiration</td> <td colspan="2">5 years</td> </tr> </table>		Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		Manufactured and Distributed by:	Tatebayashi Shokakudo Co., Ltd.		Dosage form	Powder		Packaging unit	2 g x 21 pouches	2 g x 60 pouches	Manufacturer's suggested retail price	JPY 2,404 (JPY 2,185 excluding tax)	JPY 6,072 (JPY 5,520 excluding tax)	JAN Code	4987117670448	4987117670417	Expiration	5 years
Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.																						
Manufactured and Distributed by:	Tatebayashi Shokakudo Co., Ltd.																						
Dosage form	Powder																						
Packaging unit	2 g x 21 pouches	2 g x 60 pouches																					
Manufacturer's suggested retail price	JPY 2,404 (JPY 2,185 excluding tax)	JPY 6,072 (JPY 5,520 excluding tax)																					
JAN Code	4987117670448	4987117670417																					
Expiration	5 years																						
<p>INDICATIONS</p> <p>Adiposis</p>	<p>DOSAGE AND ADMINISTRATION</p> <p>Take the medicine with water or warm water between meals.</p> <table border="1"> <thead> <tr> <th>Age</th> <th>Dose</th> <th>Daily dose</th> </tr> </thead> <tbody> <tr> <td>Adults (16 years or over)</td> <td>1 pouch</td> <td rowspan="2">3 times</td> </tr> <tr> <td>7 to 15 years</td> <td>1/2 pouch</td> </tr> <tr> <td>Under 7 years</td> <td colspan="2">Do not use.</td> </tr> </tbody> </table> <p>Between meals mean 2 to 3 hours after meals. <Precautions> For children, use under parental guidance.</p>		Age	Dose	Daily dose	Adults (16 years or over)	1 pouch	3 times	7 to 15 years	1/2 pouch	Under 7 years	Do not use.											
Age	Dose	Daily dose																					
Adults (16 years or over)	1 pouch	3 times																					
7 to 15 years	1/2 pouch																						
Under 7 years	Do not use.																						

PRECAUTIONS FOR USE

☒ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.) Lactating mothers should avoid using the medicine or stop breastfeeding.

Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using this medicine.

- (1) Persons under treatment by a physician
- (2) Pregnant women or possibly pregnant women
- (3) Persons who are physically weak (declining of strength or constitutional weakness)
- (4) Persons who have gastrointestinal weakness and are prone to diarrhea
- (5) Persons with a history of rash, redness, itching, etc. due to any drug
- (6) Persons who are taking the following drugs

Cathartic drugs (laxatives)

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected body system	Symptoms
Skin	Rash, redness, itching
Gastrointestinal	Diarrhea accompanied by severe abdominal pain, abdominal pain, constipation, feeling queasy/vomiting
Others	Oedema

3. The following symptoms may be adverse reactions to the medicine. If any of these symptoms persist or get worse, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Soft stool, diarrhea

4. If symptoms do not improve after using the medicine for about one month, consult a physician, pharmacist or registered salesperson with this leaflet.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Distributor Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>

Manufactured and Distributed by : Tatebayashi Shokakudo Co., Ltd.