


Brand name: Solmack Plus				
RISK CATEGORIES Second-class OTC drug CHARACTERISTICS Gastrointestinal medicine containing crude drugs Eight types of crude drugs are effective. (Turmeric, Glycyrrhiza, Ginseng, Coptis Rhizome, Clove, Gentian, Atractylodes Lancea Rhizome, Cinnamon Bark) + Carnitine Hydrochloride The medicine is effective for felling queasy (hangover, drunken sickness, sickness, churning of stomach, queasy, nausea), vomiting, dyspepsia, overeating (hyperphagia), overdrinking (binge drinking), and loss of appetite. It contains turmeric and glycyrrhiza, and has a cool menthol taste.				
				
ACTIVE INGREDIENTS		For further information:		
Per 2 bottles (50 mL) daily				
Turmeric Fluidextract 0.3 mL (equivalent to 300 mg of the crude drug)				
Glycyrrhiza Extract 135 mg (equivalent to 1,930.5 mg of the crude drug)				
Ginseng Fluidextract 0.5 mL (equivalent to 500 mg of the crude drug)				
Coptis Rhizome Tincture 0.826 mL (equivalent to 247.8 mg of the crude drug)				
Clove Tincture 0.126 mL (equivalent to 37.8 mg of clove)				
Gentian Tincture 0.376 mL (equivalent to 112.8 mg of the crude drug)				
Atractylodes Lancea Rhizome Fluidextract 1.2 mL (equivalent to 1,200 mg of the crude drug)				
Cinnamon Bark Tincture 0.25 mL (equivalent to 75 mg of the crude drug)				
Carnitine Hydrochloride 120 mg				
Inactive ingredients: Honey, White Soft Sugar, Acacia, Calcium Disodium Edetate, l-Menthol, Alcohol, Sodium Benzoate, pH Adjuster, Benzyl Benzoate, Flavor (≤0.8mL of alcohol/bottle)				
<ul style="list-style-type: none"> Shake well before use. As this medicine contains crud drugs, it may cause precipitation. 				
INDICATIONS				
Felling queasy (hangover, drunken sickness, sickness, churning of stomach, queasy, nausea), vomiting, dyspepsia, overeating (hyperphagia), overdrinking (binge drinking), heavy stomach, weak stomach, loss of appetite (decreased appetite), stomach/abdominal bloating, heartburn, heavy feeling in the chest				

PRECAUTIONS FOR USE

☒ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

- Do not use for a long period of time.

Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using the medicine.
 - (1) Persons under treatment by a physician
 - (2) Elderly
 - (3) Persons with oedema
 - (4) Persons diagnosed with hypertension, kidney disease or cardiac disease
2. The following serious symptoms may rarely occur after using the medicine. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician with this medicine.
Pseudoaldosteronism, myopathy (Weakness, numbness, tightness, and stiffness of limbs occur along with feelings of weakness and muscle pain, and they gradually get worse.)
3. If symptoms do not improve after using the medicine for five to six days, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27 Kandamishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>