| Brand name: TIOVITA Drink 1000  |   |                                      |   |   |
|---|---|--------------------------------------|---|---|
| <ul> <li>Brand name: HOVITA Drink 1000</li> <li>RISK CATEGORIES</li> <li>Third-class OTC drug</li> <li>CHARACTERISTICS</li> <li>I Nutrient tonic drink</li> <li>Third-class OTC drug Red Tiovita!</li> <li>It contains 1,000 mg of taurine as an active ingredient.</li> <li>It contains vitamin B1 and vitamin B6 twice and 10 times more than the conventional product (Tiovita Drink), respectively.</li> <li>Easy-to-drink "mixed fruit taste"</li> <li>It only has "68 kcal".</li> </ul> |   | ···································· | の発熱性消耗性疾患の                                |   |
| ACTIVE INGREDIENTS<br>Per 100 mL bottle<br>Taurine 1,000 mg, Inositol 50 mL,<br>Nicotinamide 20 mg, Thiamine Nitrate<br>(Vitamin B <sub>1</sub> ) 10 mg, Riboflavin Sodium<br>Phosphate (Vitamin B <sub>2</sub> ) 5 mg, Pyridoxine<br>Hydrochloride (Vitamin B <sub>6</sub> ) 50 mg,<br>Carnitine Hydrochloride 100 mg,<br>Anhydrous Caffeine 50 mg   | For further information:  |                                      |   |   |
|   | Manufactured and<br>Distributed by:   | TAIHO PHARMACEUTICAL CO., LTD.       |   |   |
|   | Dosage form   | Liquid                               |   |   |
|   | Packaging unit  | 100 mL x 1 bottle                    | 100 mL x 10<br>bottles                    | 100 mL x 50<br>bottles                    |
|   | Manufacturer's suggested retail price   | JPY 202<br>(JPY 184 excluding tax)   | JPY 2,024<br>(JPY 1,840<br>excluding tax) | JPY10,120<br>(JPY 9,200<br>excluding tax) |
| Inactive ingredients: White Soft Sugar,<br>Glucose, DL-Malic Acid, Propyl Gallate,<br>Sodium Benzoate, pH Adjuster, Vanillin,<br>Ethyl Vanillin, Flavor   | JAN Code  | 4987117437805                        | 4987117437812                             | 4987117437829                             |
|   | Expiration  | 3 years                              |   |   |
| <ul> <li>Urine may turn yellow because of vitamin B<sub>2</sub> contained in the medicine.</li> <li>It is nothing to worry about.</li> </ul>  | DOSAGE AND ADMINISTRATION<br>Adults (15 years and over): Take 1 bottle (100 mL) once daily.<br>• Do not use in children (under 15 years). |                                      |   |   |
| INDICATIONS<br>Nutritional fortification, weak constitution,<br>and supply of nutrients in the following<br>cases: physical fatigue, during and after<br>illness, loss of appetite, nutritional<br>disorders, febrile debilitating illness, during<br>pregnancy or lactation  | <ul> <li>Follow the dosin</li> </ul>  | g instruction.                       |   |   |

- Consultation
- 1. Skin rash may be adverse reactions to the medicine. If the symptom occurs, discontinue the use of the medicineimmediately, and consult a physician, pharmacist or registered salesperson with this medicine.
- 2. Diarrhea may occur after using the medicine. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.
- 3. If symptoms do not improve after using the medicine for a while, discontinue the use of the medicine, and consult aphysician, pharmacist or registered salesperson with this medicine.

## CONTACT

For inquiries, please contact the shop where you purchased the product or the following: Manufacturer and distributor Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays) Website: <u>https://www.taiho.co.jp/</u>