

Brand name: TIOVITA Drink Aibitasu

RISK CATEGORIES

Third-class OTC drug

CHARACTERISTICS

I Nutrient tonic drink

- "Blueberry flavor" which is easy to take for women
- The medicine contains vitamin B₁, vitamin B₂, and vitamin B₆ twice, three times, and 10 times more than the conventional product (Tiovita Drink), respectively.
- It contains 1,000 mg of taurine as an active ingredient.
- It only has "68 kcal".



ACTIVE INGREDIENTS

Per 100 mL bottle

Taurine 1,000 mg, Inositol 50 mg, Nicotinamide 20 mg, Thiamine Nitrate (Vitamin B₁) 10 mg, Riboflavin Sodium Phosphate (Vitamin B₂) 15 mg, Pyridoxine Hydrochloride (Vitamin B₆) 50 mg, Carnitine Hydrochloride 120 mg, Anhydrous Caffeine 50 mg

Inactive ingredients: White Soft Sugar, Glucose, DL-Malic Acid, Propyl Gallate, Sodium Benzoate, pH Adjuster, Vanillin, Ethyl Vanillin, Flavor

- Urine may turn yellow because of vitamin B₂ contained in the medicine. It is nothing to worry about.

INDICATIONS

Nutritional fortification, weak constitution, and supply of nutrients in the following cases: physical fatigue, during and after illness, loss of appetite, nutritional disorders, febrile debilitating illness, during pregnancy or lactation

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.			
Dosage form	Liquid			
Packaging unit	100 mL x 1 bottle	100 mL x 3 bottles	100 mL x 10 bottles	100 mL x 50 bottles
Manufacturer's suggested retail price	JPY 228 (JPY 207 excluding tax)	JPY 683 (JPY 621 excluding tax)	JPY 2,277 (JPY 2,070 excluding tax)	JPY 11,385 (JPY 10,350 excluding tax)
JAN Code	4987117439304	4987117439335	4987117439311	4987117439328
Expiration	3 years			

DOSAGE AND ADMINISTRATION

Adults (15 years and over): Take 1 bottle (100 mL) once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction.

PRECAUTIONS FOR USE

Consultation

1. Skin rash may be adverse reactions to the medicine. If the symptom occurs, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this medicine.
2. Diarrhea may occur after using the medicine. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.
3. If symptoms do not improve after using the medicine for a while, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>