#### Brand name: TIOVITA GOLD

## RISK CATEGORIES

Second-class OTC drug

## **CHARACTERISTICS**

#### I Nutrient tonic drink

- The medicine contains royal jelly, ginseng, andepimedium herb which are effective for severe fatigue! (For further information, please see ACTIVE INGREDIENTS.)
- It is also good for supply of nutrients at the timeof common cold!
- Easy-to-take 30 mL drink
- Sweat and mild "carrot flavor"
- It only has "34 kcal".



### **ACTIVE INGREDIENTS**

#### Per 30 mL bottle

Royal Jelly 100 mg, Ginseng Fluidextract 0.6 mL (equivalent to 600 mg of ginseng), Epimedium Herb Fluidextract 0.1 mL (equivalent to 100 mg of epimedium herb), Thiamine Nitrate (Vitamin B₁) 5 mg, Riboflavin Sodium Phosphate (Vitamin B₂) 5 mg, Pyridoxine Hydrochloride (Vitamin B₆) 5 mg, Nicotinamide 30 mg, Anhydrous Caffeine 50 mg (≤ 0.2 mL of alcohol)

Inactive ingredients: Citric Acid, , Propyl Gallate, Alcohol, Sodium Benzoate, pH Adjuster, Vanillin, Corn Starch, Flavor

- (1) Shake well before use. As this medicinecontains crud drugs, it may cause precipitation.
- (2) Urine may turn yellow because of vitamin B2 contained in the medicine. It is nothing to worry about.

#### **INDICATIONS**

Supply of nutrients in the following cases: febrile debilitating illness, physical fatigue, during and after illness, loss of appetite, nutritional disorders, during pregnancy or lactation, and nutritional fortification, weak constitution

## For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form	Liquid		
Packaging unit	30 mL x 1 bottle	30 mL x 3 bottles	30 mL x 10 bottles
Manufacturer's suggested retail price	JPY 443 (JPY 403 excluding tax)	JPY 1,329 (JPY 1,208 excluding tax)	JPY 4,428 (JPY 4,025 excluding tax)
JAN Code	49684315	4987117437218	4987117437355
Expiration	3 years		

## DOSAGE AND ADMINISTRATION

Adults (15 years and over): Take 1 bottle (30 mL) once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction.

# PRECAUTIONS FOR USE

## Consultation

- 1. Diarrhea may occur after using the medicine. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.
- 2. If symptoms do not improve after using the medicine for a while, discontinue the use of the medicine, and consult aphysician, pharmacist or registered salesperson with this medicine.

## CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <a href="https://www.taiho.co.jp/">https://www.taiho.co.jp/</a>