かゆみ 保湿	ила
For further information:	
Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.
Manufactured and Distributed by:	OKAYAMA TAIHO PHARMACEUTICAL CO., LTD.
Dosage form	Cream
Packaging unit	80 g
Manufacturer's suggested retail price	JPY 2,024 (JPY 1,840 excluding tax)
JAN Code	4987117140125
Expiration	3 years
INDICATIONS Dry skin accompanied by itching (xeroderma in the elderly/adults, dry skin in children) Precautions> (1) Be careful not to get the medicine into the eyes medicine gets into the eyes, immediately wash with lukewarm water. If symptoms are severe, consult a ophthalmologist. (2) For children, use under parental guidance. (3) For external use only. (4) Clean the affected area and surrounding skin brows. (5) Follow the prescribed dosing instructions. (6) As this is not a cosmetics, use only for the affected in the indications, and do not use on the face purpose of basic skin care, etc.	
	For further informatio Distributed by: Manufactured and Distributed by: Dosage form Packaging unit Manufacturer's suggested retail price JAN Code Expiration DOSAGE AND ADM Apply an appropriate several times daily. <precautions> (1) Be careful not to g medicine gets into the lukewarm water. If sy ophthalmologist. (2) For children, use (3) For external use of (4) Clean the affected (5) Follow the prescri (6) As this is not a co listed in the indication</precautions>

PRECAUTIONS FOR USE X The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

Do not apply to the following areas.

(1) Around the eyes, mucous membranes (e.g. mouth, nasal cavity, vagina, etc.)

(2) Inflammation or wounds

(3) Severe skin sore or cracking of the skin

Consultation

1. The following persons should consult a physician, pharmacist or registered distributor before using the medicine.

(1) Persons under treatment by a physician

(2) Persons with a history of allergic symptoms (e.g. rash, redness, itching, contact dermatitis, etc.) or irritation symptoms

due to any drug

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered distributor with this leaflet.

Affected body system	Symptoms
Skin	Contact dermatitis, rash, redness, itching, irritation (pain, feeling hot, tingling sensation), swelling, scab-like skin exfoliation
	using the medicine for about two weeks, discontinue the use of the medicine, and egistered distributor with this leaflet.
Distributor Contact: Customer Re -27 Kandanishiki-cho, Chiyoda-	hours: 9:00-17:00 (Excluding weekends and holidays)